

ESM 2. Clinical outcomes at 12 months – additional data

	All patients	Critical limb ischaemia	No critical limb ischaemia	p-value
Patients	357	60	297	
<i>Rutherford category</i>				
0	115 (37.0)	16 (32.7)	99 (37.8)	<0.001
1	89 (28.6)	8 (16.3)	81 (30.9)	
2	36 (11.6)	2 (4.1)	34 (13.0)	
3	18 (5.8)	1 (2.0)	17 (6.5)	
4	10 (3.2)	4 (8.2)	6 (2.3)	
5	4 (1.3)	3 (6.1)	1 (0.4)	
6	2 (0.6)	2 (4.1)	0 (0.0)	
Unknown or impacted*	37 (11.9)	13 (26.5)	24 (9.2)	
TVR (recurrent PTA, lysis)	43 (13.8)	9 (18.4)	34 (13.0)	0.316
Non-target vessel revascularisation	23 (7.4)	7 (14.3)	16 (6.1)	0.045
Major amputations, target and non-target leg	8 (2.6)	5 (10.2)	3 (1.1)	<0.001
Follow-ups relative to mortality**	313 (87.7)	50 (83.3)	263 (88.6)	0.262
Follow-up duration – months (including time to event)	11.4 ± 3.3	10.1 ± 4.3	11.7 ± 3.0	0.001
All thrombotic events including peri-procedural events	17 (5.5)	3 (6.1)	14 (5.3)	0.826
<i>Death</i>				
Not related to intervention	17 (5.5)	6 (12.2)	11 (4.2)	<0.001
Possibly related to Intervention	2 (0.6)	1 (2.0)	1 (0.4)	
Cardiac death	3 (1.0)	3 (6.1)	0 (0.0)	

Data are n (%) or mean \pm standard deviation. For continuous variables p-values are from unpaired student t-test and chi square for categorical variables. TVR = target vessel revascularization; PTA = Percutaneous transluminal angioplasty.

*presence of claudication due to contralateral leg, non-vascular factors or amputations

**follow-ups by telephone