

ESM 3. Impact of different variables (derived from the literature) on freedom from clinically-driven target lesion revascularisation in a logistic regression model

Variable	OR (95% CI)	p-value
Age – years	0.990 (0.945 – 1.039)	0.692
Male gender	1.981 (0.797 – 4.928)	0.141
Diabetes mellitus – yes/no	2.384 (0.954 – 5.958)	0.063
Hypertension – yes/no	0.498 (0.128 – 1.929)	0.313
Hypercholesteremia – yes/no	0.995 (0.383 – 2.579)	0.991
Current smoking – yes/no	0.808 (0.461 – 1.414)	0.455
Critical limb ischaemia – yes/no	1.304 (0.358 – 4.745)	0.687
TASC II C/D lesion – yes/no	1.196 (0.435 – 3.286)	0.729
No distal run-off (no vessel) – yes/no	2.733 (0.901 – 8.290)	0.076
Vessel diameter – mm	1.027 (0.647 – 1.630)	0.911
Total lesion length – cm	0.998 (0.993 – 1.004)	0.600
Chronic total occlusion – yes/no	1.764 (0.705 – 4.417)	0.225
Predilation DCB – yes/no	0.468 (0.184 – 1.189)	0.110

OR = odds ratio; CI = confidence interval.