

Electronic supplementary material

ESM 1. Comparison of the cardiovascular risk factors in the various vascular beds affected by atherosclerosis

	PAD	CS	CAD	No
	N=318	N=159	N=2572	atherosclerosis
				N=705
Age (years)	65.4(8.77)	67.3(8.13)	63.8(9.86)	58.4 (12.1)
Female sex (%)	18,6	20,8	25,3	48,8
BMI (kg/m ²)	27.2(4.11)	27.2(4)	27.5(4.01)	27.4 (4.31)
LDL-C (mg/dL)	111(29.6)	110(34.2)	116(34.8)	120 (32.3)
HDL-C (mg/dL)	36(10.6)	37(10.3)	37.6(10.2)	42.8 (11.8)
TG (mg/dL)	164(121-223)	153(113-204)	150(113-202)	133 (97–194)
FGLUC (mg/dL)	109(96.8-141)	104(95-127)	104(94.6-122)	98.8 (91.5–109)
HbA1c (%)	6.82(1.64)	6.55(1.43)	6.4(1.3)	6 (1)
SBP (mmHg)	149(25.5)	148(24.2)	143(23.7)	136 (22.1)
DBP (mmHg)	80.7(12.2)	80.5(11.6)	81.2(11.5)	80.3 (11.3)
PP (mmHg)	73.4(20.2)	72.8(19.7)	65.9(18.5)	59.2 (17.1)
eGFR (ml/min/1.73 m ²)	75.6(21.2)	75.6(22.1)	80.6(20.2)	86.1 (19.1)
Alcohol intake (g eth/d)	2.68(0-24)	2.8(0-24)	3(0-24)	3.36 (0–24)
hsCRP (mg/L)	4.32(2.06-9.98)	3.95(1.49-7.64)	3.84(1.5-9.24)	2.12 (0.98–5.61)
NT-proBNP (pg/mL)	420(152-1610)	432(177-1330)	333(126-967)	157 (70-492)
Hypertension (%)	83,6	83	75,3	62,4
T1DM (%)	1,89	0,629	0,35	0
T2DM (%)	55,7	43,4	43,3	26,5
Current smoker (%)	32,1	17	24	20,4
Ex-smoker (%)	50	57,2	44,9	27,8
CAD (%)	94,7	93,1	100	0
PAD (%)	100	34	11,7	0
CS (%)	17	100	5,75	0
1-vessel CAD (%)	19,7	20,4	37,2	0
2-vessel CAD (%)	26,1	26,8	37,4	0
3-vessel CAD (%)	53,5	54,8	38,5	0

Notes. PAD: peripheral arterial disease, CS: carotid stenosis, CAD: coronary artery disease.