

Electronic supplementary material

ESM 1. Depressive symptoms and mean performances on neuropsychological measures at pre and post of patients with at least one comorbid anxiety disorder versus those without

	Pre		Statistics, pre, and between-group <i>t</i> -tests	Post		Statistics, post, between-group <i>t</i> -tests, and repeated measures ANCOVAs
	MDD (<i>n</i> = 23) <i>M</i> (<i>SD</i>)	MDDA (<i>n</i> = 22) <i>M</i> (<i>SD</i>)		MDD (<i>n</i> = 23) <i>M</i> (<i>SD</i>)	MDDA (<i>n</i> = 22) <i>M</i> (<i>SD</i>)	
<i>Depression</i>						
HDRS	19.26 (4.84)	22.50 (6.67)	$t(38.19) = 1.87, p = .07, d = 0.56$	8.83 (7.09)	10.59 (7.89)	$t(43) = 0.79, p = .43, d = 0.24$
BDI	22.22 (5.70)	26.55 (9.78)	$t(33.47) = 1.83, p = .08, d = 0.54$	12.48 (9.05)	15.65 (9.72)	$t(43) = 1.13, p = .26, d = 0.34$
<i>Neuropsychological measures</i>						
Test d2, concentration performance index	160.04 (40.82)	166.52 ^a (47.24)		189.48 (45.99)	185.29 (54.81)	Group: $p = .70, \eta_p^2 < .001$ Time: $p < .001, \eta_p^2 = .26$ Group x Time: $p = .09, \eta_p^2 = .07$
Trail Making Test A, sec.	30.64 (17.39)	25.78 (11.65)		22.00 (7.28)	23.73 (10.43)	Group: $p = .22, \eta_p^2 = .04$ Time: $p = .35, \eta_p^2 = .02$ Group x Time: $p = .017, \eta_p^2 = .13$
Trail Making Test B, sec.	75.70 (35.64)	69.78 ^a (35.72)		56.61 (25.65)	56.20 (27.89)	Group: $p = .45, \eta_p^2 = .01$ Time: $p = .36, \eta_p^2 = .02$ Group x Time: $p = .40, \eta_p^2 = .02$

Note. MDD = depression without comorbid anxiety; MMDA = depression with comorbid anxiety; HDRS = Hamilton Rating Scale for Depression; BDI = Beck Depression Inventory; ^a *n* = 21.