

Tabelle E5. Post-hoc Vergleiche für die Fehlerkategorien im Rahmen der Messwiederholungs-ANOVA im Zwei-Gruppen-Fall

Paarweiser Vergleich der Fehlerkategorien		Mittlere Differenz	Standardfehler	<i>p</i>
A	H	7.49	0.42	< .0001
	LV	2.15	0.39	< .0001
	U	7.96	0.43	< .0001
	SW	8.02	0.40	< .0001
	GK	-9.76	0.61	< .0001
	LF	-3.06	0.40	< .0001
	AW	8.11	0.43	< .0001
H	LV	-5.34	0.30	< .0001
	U	0.47	0.12	.004
	SW	0.53	0.18	.121
	GK	-17.25	0.42	<.0001
	LF	-10.55	0.28	<.0001
	AW	0.62	0.15	.002
LV	U	5.81	0.31	<.0001
	SW	5.87	0.32	<.0001
	GK	-11.91	0.50	<.0001
	LF	-5.21	0.30	<.0001
	AW	5.96	0.32	<.0001
U	SW	0.06	0.18	1.00
	GK	-17.72	0.43	<.0001
	LF	-11.02	0.28	<.0001
	AW	0.15	0.15	1.00
SW	GK	-17.78	0.48	<.0001
	LF	-11.08	0.33	<.0001
	AW	0.09	0.18	1.00
GK	LF	6.70	0.46	<.0001
	AW	17.87	0.46	<.0001
LF	AW	11.17	0.30	<.0001

Anmerkungen. Unter Berücksichtigung der Bonferroni-Korrektur sind p-Werte < .0018 statistisch signifikant.