

Table. *Previously published factor solutions for the SPAI-C*

Publication	Country	Sample Characteristics	Method	Factors and Items Included	Excluded Items
Beidel et al. (1995)	USA	Combined Sample (Clinical and Control) N = 154 8-17 years	PCA	<i>Assertiveness/General Conversation</i> : 2, 8, 9, 10, 11, 12, 13, 14, 15, 18, 19, 21, 24 <i>Traditional Social Encounters</i> : 6, 9, 17, 19, 20, 21, 24, 25, 26 <i>Public Performance</i> : 1, 2, 3, 4, 5, 16, 17	7, 22, 23
Beidel et al. (1996)*	USA	Combined Sample (Clinical and Non-clinical)	PCA	<i>Assertiveness</i> : 8, 10, 11, 12, 13, 17, 18 <i>General Conversation</i> : 1, 3, 7, 14, 15, 16 <i>Physical and Cognitive Symptoms</i> : 21, 24, 25, 26 <i>Avoidance</i> : 2, 6, 19, 20 <i>Public Performance</i> : 3, 4, 5, 16	9, 22, 23
Gauer et al. (2005)	Brazil	Community Sample N = 1,871 9-14 years	Factor analysis with varimax rotation	<i>Assertiveness</i> : 2, 10, 11, 12, 13, 18 <i>Avoidance/Social Encounters</i> : 6, 7, 9, 14, 15, 19, 20 <i>Public Performance</i> : 1, 3, 4, 5, 8, 16, 17 <i>Physical and Cognitive Symptoms</i> : 21, 22, 24, 25, 26	23
Aune et al. (2008)	Norway	Community Sample N = 1,439 11-14 years	EFA and CFA	<i>Assertiveness</i> : 10, 11, 12, 13, 18 <i>Physical/Cognitive Symptoms</i> : 21, 22, 25, 26 <i>Public Performance</i> : 2, 3, 4, 5, 8, 16 <i>Social Encounters</i> : 6, 7, 9 <i>Avoidance</i> : 19, 20	1, 14, 15, 17, 23, 24
Olivares et al. (2010)	Spain	Community Sample N = 1,588 10-17 years	CFA	<i>Public Performance</i> : 1, 2, 3, 4, 5, 8, 16 <i>Assertiveness</i> : 2, 10, 11, 12, 13, 17, 18, 23 <i>Fear and Avoidance/Escape in Social Encounters</i> : 6, 7, 9, 14, 15, 19, 20, 22 <i>Cognitive and Psychophysiological Interferences</i> : 21, 22, 24, 25, 26	n/a
Ogliari et al. (2012)**	Italy	Community Sample N = 228 8-11 years	CFA	<i>Assertiveness</i> : 8, 10, 11, 12, 13, 17, 18 <i>General Conversation</i> : 1, 7, 14, 15 <i>Physical/Cognitive Symptoms</i> : 21, 24, 25, 26 <i>Avoidance</i> : 6, 10, 19, 20 <i>Public Performance</i> : 3, 4, 5, 16 A hierarchical model with 5 first-order factors in which the 5 factors were predicted by a second-order social phobia factor was also tested.	2, 9, 22, 23
Cederlund & Öst (2013)	Sweden	Clinical Sample N = 59 8-14 years	EFA	<i>Fear of Social Interactions/Social Encounters</i> : 9, 10, 11, 12, 13, 14, 15, 18, 19 <i>Public Performance</i> : 2, 4, 5, 8, 17 <i>Symptoms</i> : 1, 7, 21, 22, 23, 24, 25, 26	3, 6, 16, 20

Notes: *Storch et al. (2004) also reported that the Beidel et al. (1996) five-factor solution was an adequate fit for their data from a community sample of 13-17 year olds in the USA using CFA ($N = 1,147$) **Hierarchical model proposed by Ogliari et al. (2012).